

Nuclear Reaction Rates in Dense Plasmas

V. I. Savchenko

⁺ *Princeton University, PPPL, Forrestal Campus, Princeton, N. J. 08543*

We solve the quasiclassical problem of tunneling through an external potential barrier in a dense plasma, where the tunneling particles undergo simultaneous collisions with other particles in thermodynamic equilibrium. Under such conditions the spectral density of states available to the particle has a Lorentz shape, rather than the delta function, which leads to a quantum tail in the particle momentum distribution function. We show that, this tail indeed significantly alters the average nuclear reaction rates, which supports earlier suggestions. This rate can be many orders of magnitude higher than would be normally calculated by averaging over the Maxwell distribution of energies.

The particle distribution over momenta, $f(\mathbf{p})$, acquires non-Maxwellian tail in thermodynamic equilibrium [1] due to quantum effects, while the particle distribution over energies, $\tilde{f}(\omega)$, remains Maxwellian. An interesting suggestion was made in Ref. [2], that the rates of various processes, including nuclear reactions, may be significantly increased by the quantum tail of the *momentum* distribution. A final formula for the ionization rate in a closed form was obtained in [2], by using the Born approximation. The formula for the nuclear reaction rates was found in [3] under assumption that it is correct to substitute the particle momentum in the form $\epsilon_{\mathbf{p}} = p^2/2m$ into the known quasiclassical cross-section $\sigma(\omega)$ and then average it over the particle distribution over momenta $f(\mathbf{p})$ rather than the distribution over energies $\tilde{f}(\omega)$. This procedure becomes even more unclear, if we take into account the fact, that the momentum, $\mathbf{p}$, and energy, ω , of a *colliding* particle are *independent* variables not connected by the usual dispersion relation $\delta(\omega - \mathbf{p}^2/2m)$ [1,4,2]. Note that the averaging of $\sigma(\omega)$ over the distribution over energies $\tilde{f}(\omega)$, which is Maxwellian would give the usual reaction rates [5] rather than the rates found in [3], which are accelerated by many orders of magnitude in certain regimes. It is therefore important to find the nuclear reaction rates from first principles.

In this paper we rigorously solve the quasiclassical problem of tunneling through the potential barrier, when the tunneling particles undergo simultaneous collisions with other particles of the plasma maintained in thermodynamic equilibrium, (see Fig. 1). The plasma has density and

temperature n_0, T and is fully ionized. We use the Green-function technique [6–8], and do not rely on any assumptions [3] about any averaging procedure. We show that, if the requirements of the quasiclassical approximation are fulfilled with respect to both the barrier and collisions, our final result for the nuclear reaction rate supports the idea of averaging over momentum postulated in [3].




FIG. 1.

According to the diagram technique of Keldsyh and Korenman [7,8] the kinetic properties of the particles are described by the Green function $G^{-+}(X_1, X_2)$; we adopt the notation $X = (t, \mathbf{x})$. Kinetic equations for this function [7,8] include the interaction between the particles as well as their interaction with the external field. Therefore, the rate of change of this function, evaluated at any point after the barrier, will give the tunneling rate $K(\mathbf{x}, t)$ for the particles colliding simultaneously with other particles:

$$K(\mathbf{x}, t) = \left[(\partial_{t_1} - \partial_{t_2}) G^{-+}(t_1 \mathbf{x}_1; t_2 \mathbf{x}_2) \right]_{X_1 \rightarrow X_2 = X}$$

$$= 2 \left[\partial_{t_1} G^{-+}(t_1 \mathbf{x}_1; t_2 \mathbf{x}_2) \right]_{X_1 \rightarrow X_2 = X}, \quad (1)$$

In the second line of (1) we anticipated that, G^{-+} depends only on $t_1 - t_2$.

Now we will proceed with solution of the kinetic equations for $G^{-+}(X_1, X_2)$. They can be written as Dyson equations for $G_{12}^{\alpha\beta}$ [6]:

$$G_{12}^{\alpha\beta} = G_{12}^{(0)\alpha\beta} + \int G_{14}^{(0)\alpha\gamma} \Sigma_{43}^{\gamma\delta} G_{32}^{\delta\beta} d^4 X_4 d^4 X_3, \quad (2)$$

where we use subscripts 1..4 to denote dependence on $X_1 \dots X_4$. Dependence on $G_{12}^{(0)\alpha\beta}$ can be eliminated [6] by acting on both sides of the Eq. (2) with the operator

$$\hat{G}_1^{-1} = i \frac{\partial}{\partial t_1} + \frac{\Delta_1}{2m} - U(z_1) \equiv i \frac{\partial}{\partial t_1} + \frac{\Delta_1^\perp}{2m} + \hat{L}(z_1),$$

where $\Delta_1^\perp \equiv \partial_{x_1}^2 + \partial_{y_1}^2$. Since we are interested in the steady state solution, one can see that, the properties of this operator can be fully accounted for if we introduce the function $g(z, k)$, such that

$$\left[\frac{1}{2m} \frac{\partial^2}{\partial z^2} - U(z) \right] g(z, k) = -\epsilon_k g(z, k), \quad (3)$$

where $\epsilon_k = k^2/2m$. Since the operator $\hat{L}$ is Hermitian, we will use the property of its eigenfunctions $g(z, k)$:

$$\int_{-\infty}^{\infty} g(z, k_1) g^*(z, k_2) dz = 2\pi \delta(k_1 - k_2) \quad (4)$$

Now let us solve for $G_{12}^{(0)-+}$, which satisfies

$$\hat{G}_1^{-1} G_{12}^{(0)-+} = 0. \quad (5)$$

It is easier to find $G_{12}^{(0)-+}$ by using its definition

$$G_{12}^{(0)-+} = i \langle \hat{\psi}_2^\dagger \hat{\psi}_1 \rangle, \quad (6)$$

where $\hat{\psi}^\dagger, \hat{\psi}$ are the Heisenberg operators of creation and annihilation of the particles at a point X , and $\langle \dots \rangle$ means quantum and statistical averaging. Since $\hat{\psi}$ evolves according to $\hat{G}^{-1} \hat{\psi} = 0$, one can see, that it is equal to

$$\hat{\psi}(X) = \sum_{\mathbf{q}_\perp k} \hat{a}_{\mathbf{q}_\perp k} g(z, k) e^{-i\epsilon_{\mathbf{q}_\perp k} t + i\mathbf{q}_\perp \cdot \mathbf{x}^\perp} \quad (7)$$

Here $\hat{a}_{\mathbf{q}_\perp k}$ is the annihilation operator of the particle with momentum $(\mathbf{q}_\perp, k)$, and $\epsilon_{\mathbf{q}_\perp k} \equiv \mathbf{q}_\perp^2/2m + k^2/2m$.

We substitute $\psi(X)$ from (7) into Eq. (6) and obtain

$$G_{\omega \mathbf{q}_\perp}^{(0)-+}(z_1, z_2) = 2\pi i \int_{-\infty}^{\infty} \frac{dk}{2\pi}$$

$$\delta(\omega - \epsilon_{\mathbf{q}_\perp k} + \mu) n(\mathbf{q}_\perp k) g(z_1, k) g^*(z_2, k), \quad (8)$$

where we denote $\langle a_{\mathbf{q}_\perp k}^\dagger a_{\mathbf{q}_\perp k} \rangle \equiv n(\mathbf{q}_\perp k)$; μ is the chemical potential.

It is easy to see that, in thermodynamic equilibrium we are considering, the function $n(\mathbf{q}_\perp k)$ is equal to Fermi distribution $n_F(\epsilon_{\mathbf{q}_\perp k})$. To convince ourselves we take $G_{\omega \mathbf{q}_\perp}^{(0)-+}(z_1, z_2)$ at such values of z_1, z_2 , when $U(z)$ is negligible and $g(z, k) \Rightarrow \exp(ikz)$. One more Fourier transform $\int \exp[-iq(z_1 - z_2)] d(z_1 - z_2)$ will give $G_{\omega \mathbf{q}}^{(0)-+}$, which has to coincide with the known expression [6] $G_{\omega \mathbf{q}}^{(0)-+} = 2\pi i \delta(\omega - \epsilon_{\mathbf{q}} + \mu) n_F(\epsilon_{\mathbf{q}})$.

Before proceeding further with solving for $G^{-+}(X_1, X_2)$ we first find $K^{(0)}$, which is the tunneling rate obtained by using $G_{12}^{(0)-+}$ in Eq. (1). Since we are interested in $K^{(0)}$ behind the barrier, where $U(z) = 0$, we can exchange ∂_{t_1} for $i\Delta_1/2m$ as is clear from the form of the operator $\hat{G}_1^{-1}$ and Eq. (5). Then the formula for the rate $K^{(0)}$ takes the form:

$$K^{(0)}(z) = \frac{i}{2m} \int \frac{d\omega}{2\pi} \int \frac{d^2 \mathbf{q}_\perp}{(2\pi)^2} \left[\frac{\partial^2}{\partial \zeta^2} G_{\omega \mathbf{q}_\perp}^{(0)-+}(z, \zeta) \right]_{\zeta=0}, \quad (9)$$

where $z = (z_1 + z_2)/2$, $\zeta = z_1 - z_2$.

In deriving (9) we expressed $G_{12}^{(0)-+}$ through its Fourier transform $G_{\omega \mathbf{q}_\perp}^{(0)-+}(z, \zeta)$ with respect to “fast” space $\mathbf{r}_\perp = \mathbf{r}_1^\perp - \mathbf{r}_2^\perp$ and time $\tau = t_1 - t_2$ variables and took the limit $\mathbf{r}, \tau \rightarrow 0$ as prescribed by Eq. (1).

Finally we need to know the function $g(z, k)$ in the region behind the barrier. We will use the standard quasiclassical expression [9]

$$g(z, k) = \frac{C_k}{\tilde{q}(z, k)^{1/2}} \exp \left[i \int_{z_*}^z \tilde{q}(z', k) dz' \right] \quad (10)$$

$$C_k = A_k \exp \left[- \int_{z_a}^{z_b(k)} \tilde{q}(z', k) dz' \right] \quad (11)$$

$$\tilde{q}(z, k) = \sqrt{2m(\epsilon_k - U(z))} \quad (12)$$

The square of the exponential factor in Eq. (11) is the tunneling coefficient, $\tilde{W}(k)$ [10]. We multiply it by a factor $A_k = (S(\epsilon_k)/\epsilon_k)^{1/2}$, which takes into account nuclear physics effects, not considered here. $S(\epsilon_k)$ is the astrophysical factor [11].

To find $K^{(0)}$ from Eq. (9) we substitute Eqs. (10)-(12) in Eq. (8), use $z_1 = z + \zeta/2$, $z_2 = z - \zeta/2$ and take ∂_ζ^2 derivative in the quasi-classical sense keeping only zero-th order terms. The result of the differentiation is

$$\left[\frac{\partial^2}{\partial \zeta^2} [g(z_1, k)g^*(z_2, k)] \right]_{\zeta=0} = -\tilde{q}(z, k)C_k^2 \quad (13)$$

Note, that $\tilde{q}(z, k) = k$ is true behind the barrier.

The problem of tunneling collision between particles m_1 and m_2 can be reduced to the one-dimensional motion of a particle with reduced mass $m_r = m_1 m_2 / (m_1 + m_2)$ in the external potential $U(r)$ [9]. We make this reduction in the original Hamiltonian and then second quantize it. Then we use spherical coordinates, assume spherical symmetry of the tunneling collision and perform calculations similar to those explained above. The final result for $K^{(0)}$ is

$$K^{(0)} = n_0 \int_0^\infty \frac{4\pi k^2 dk}{(2\pi)^3} \bar{n}_M(k) \frac{k}{m_r} \frac{S(\bar{\epsilon}_k)}{\bar{\epsilon}_k} W(k), \quad (14)$$

where $\bar{\epsilon}_k = k^2/2m_r$. $W(k)$ is obtained from $\tilde{W}(k)$ by $z \rightarrow r$, $\bar{n}_M(k)$ is the Maxwell distribution which depends on m_r .

It agrees with the answer obtained in [5] by averaging the quasiclassical tunneling factor $W(k)$ over the Maxwell distribution, $\langle n_M(v)\sigma(m_r v^2/2)v \rangle$. Hence, we see, that the dispersion relation $\delta(\omega - \epsilon_{\mathbf{q}})$ was implicitly used in [5], which corresponds to an approximation of instantaneous, two-body collisions. Now we will not make this assumption and proceed with finding the tunneling rate, K , of the particles, which always collide and hence are never “in between collisions”.

By using [6,4], we find

$$\left(\frac{q}{m} \frac{\partial}{\partial z} + U'(z) \frac{\partial}{\partial q} \right) G_{\omega \mathbf{q}}^{-+}(z) = -\gamma_{\omega \mathbf{q}}(z) G_{\omega \mathbf{q}}^{-+}(z) - \gamma_{\omega \mathbf{q}}(z) n_F(\omega) (G_{\omega \mathbf{q}}^R(z) - G_{\omega \mathbf{q}}^A(z)), \quad (15)$$

where we denote $\Sigma^R - \Sigma^A \equiv i\gamma_{\omega \mathbf{q}}(z) \equiv i\gamma_{\omega \mathbf{q}\perp}(q, z)$. Note, that $\gamma_{\omega \mathbf{q}}(z)$ is a non-linear function of $G^{\alpha\beta}$ [4,6]. We now analyze Eq. (15) qualitatively, which will help us to find that part of the solution, which makes the largest contribution to K .

In the region *away* from the barrier we can neglect the LHS of Eq. (15), and obtain:

$$\tilde{G}_{\omega \mathbf{q}}^{-+}(z) = -n_F(\omega) (G_{\omega \mathbf{q}}^R(z) - G_{\omega \mathbf{q}}^A(z)) \quad (16)$$

Therefore, Eq. (15) will allow us to propagate this solution into the region *behind* the barrier.

We can find the first integral and formally integrate Eq. (15) along trajectories, which will lead to a sum of solutions of homogeneous and “inhomogeneous” equations (due to the last term in (15)). As can be seen from Eq. (15), “inhomogeneous” solution will involve the integral of the product of $\gamma_{\omega \mathbf{q}}(z)$ and $G_{\omega \mathbf{q}}^R(z)$ or $G^{R*} = G^A$. Qualitatively, it means, that in the region of z behind the barrier it will be proportional to a product of at least two exponential factors $W(k)$. This is so, because both $\gamma_{\omega \mathbf{q}}(z)$ and $G_{\omega \mathbf{q}}^R(z)$ will depend on the integral of the product $g(z_1, k)g^*(z_2, k)$ and such product is $\propto W(k)$ as can be seen from Eqs. (10)-(12).

It is obvious that, the solution of the “homogeneous” equation, G_f^{-+} , involves only one factor of $W(k)$, since $G_f^{-+} \propto gg^*$, see Eq. (15). Therefore, we will find only the solution to the “homogeneous” equation, which we will use to obtain K . We supplement this homogeneous equation with the boundary condition, $\tilde{G}^{-+}$ from Eq. (16) imposed at $z = z_*$ away from the barrier, see Fig. 1.

Since the largest contribution to K is made by G_f^{-+} one can see, that we can use Eq. (9) to find K if we substitute G_f^{-+} instead of $G^{(0)-+}$ in (9). Therefore, we find G_f^{-+} with dependence

on z_1, z_2 from the very beginning by applying the transformation

$$\int \int g(z_1, k_1) g^*(z_2, k_2) \frac{dk_1}{2\pi} \frac{dk_2}{2\pi} \quad (17)$$

to Eq. (2). We use the boundary condition, Eq. (16) and obtain the following answer:

$$G_f^{-+} = 2\pi i n_F(\omega) \int g_k(z_1) \delta_\gamma(\omega - \epsilon_{\mathbf{q}_\perp k}) g_k^*(z_2) \frac{dk}{2\pi} \quad (18)$$

where $g_k(z) \equiv g(z, k)$ and we used the well-known equilibrium solution [4]

$$G_{\omega \mathbf{q}_\perp}^R(k) - G_{\omega \mathbf{q}_\perp}^A(k) = \frac{\gamma_{\omega \mathbf{q}_\perp}(k)}{(\omega - \epsilon_{\mathbf{q}_\perp k})^2 + (\gamma_{\omega \mathbf{q}_\perp}(k))^2}$$

$$G_{\omega \mathbf{q}_\perp}^R(k) - G_{\omega \mathbf{q}_\perp}^A(k) \equiv \delta_\gamma(\omega - \epsilon_{\mathbf{q}_\perp k}) \quad (19)$$

Comparing Eq. (8) with (18), (19) we see, that the effect of collisions, which makes the largest contribution to K can be described as $\delta(\omega - \epsilon_{\mathbf{q}_\perp k}) \Rightarrow \delta_\gamma(\omega - \epsilon_{\mathbf{q}_\perp k})$, as can be expected on the intuitive grounds. Note also, that we can obtain the answer (18), (19) by following the same steps leading to Eq. (8) if we make a substitution $\epsilon_{\mathbf{q}_\perp k} \rightarrow \epsilon_{\mathbf{q}_\perp k} + i\gamma_{\mathbf{q}_\perp k}$. This means that, we have to use *damped* oscillators instead of *undamped* ones for second quantization of *colliding* particles.

Now we substitute Eqs. (18), (19) into (9) and integrate $\int d\omega/2\pi$ by using the result of [1]:

$$\int \frac{d\omega}{2\pi} n_M(\omega) \delta_\gamma(\omega - \epsilon_{\mathbf{q}_\perp k}) = n_M(\epsilon_{\mathbf{q}_\perp k}) + \delta n_\gamma(\mathbf{q}_\perp k), \quad (20)$$

where $\delta n_\gamma(\mathbf{q}_\perp k)$ is the “quantum tail”.

At this point we are ready to perform the same steps as those leading to Eq. (14) in order to describe the tunneling of particles m_1 and m_2 . The answer for K is:

$$K = n_0 \int_0^\infty \frac{4\pi p^2 dp}{(2\pi)^3} \frac{S(\bar{\epsilon}_p)}{\bar{\epsilon}_p} \quad (21)$$

We now evaluate Σ^R through collision frequency, ν_T , as in [1,12], carry out the integral in (21) and form the ratio $r_{12} = K_\gamma/K_M$:

$$r_{12} = \frac{3^{19/2}}{8\pi^{3/2}} \sum_j \frac{h\nu_T(m_{coll})}{T} \left(\frac{m_{coll}}{m_r} \right)^{7/2} \frac{e^{\tau_{12}}}{\tau_{12}^8} \quad (22)$$

Here $m_{coll} = m_r m_j / (m_r + m_j)$, $m_r = m_1 m_2 / (m_1 + m_2)$, $\tau_{12} = 3(\pi/2)^{2/3} \left(\frac{100 Z_1^2 Z_2^2 A_{12}}{T_k} \right)^{1/3}$, $A_{12} = A_1 A_2 / (A_1 + A_2)$, T_k is the temperature in keV; A_1, A_2 are the atomic numbers of tunneling particles, m_j is the mass of the background particles colliding with tunneling particles. Summation is over all species of colliding background particles, including tunneling species.

This result, (21), (22), supports the idea of averaging of σv over momentum distribution postulated in [3], but differs from the answer of [3] by a factor $(m_{coll}/m_r)^3$ under the summation sign.

Consider, for example, a DT plasma of density $\rho = 10 \text{ g/cm}^3$ and temperature $T = 0.1 \text{ keV}$. Then Eq. (22) will lead to the rate accelerated by five orders of magnitude as compared to the conventional answer. We find the ratio $r_{DT} = K_{DT}^\gamma / K_{DT}^M = 8 \cdot 10^4$, with $K_{DT}^M = 2.7 \cdot 10^{-30} \text{ cm}^3/\text{sec}$, $K_{DT}^\gamma = 2.1 \cdot 10^{-25} \text{ cm}^3/\text{sec}$. In case of heavier elements and not very small $h\nu_T/T$, ratio r_{ij} becomes much higher [3].

I would like to acknowledge helpful discussions with N. J. Fisch. This work was supported by NSF-DOE grant DE-FG02-97ER54436.

-
- [1] V. M. Galitski and V. V. Yakimets, ZhETF **51**, 957 (1966).
 - [2] A. N. Starostin and N. L. Aleksandrov, Phys. Plasmas **5**, 2127 (1998); N. L. Aleksandrov and A. N. Starostin, ZhETF **113**, 1661 (1998).
 - [3] A. N. Starostin, V. I. Savchenko, N. J. Fisch submitted to Phys. Rev. Lett.
 - [4] L. P. Kadanoff and G. Baym, *Quantum Statistical Mechanics* (W. A. Benjamin, New York, 1962).
 - [5] G. Gamow and E. Teller, Phys. Rev. **53**, 608 (1938).
 - [6] E. M. Lifshitz and L. P. Pitaevskii, *Physical Kinetics* (Pergamon Press, New York, 1977).
 - [7] L. V. Keldish, Sov. Phys. JETP **20**, 1018 (1965).
 - [8] V. Korenman, Ann. Phys. **39**, 72 (1966).
 - [9] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Pergamon Press, New York, 1977).
 - [10] E. M. Lifshitz and L. P. Pitaevskii, *Relativistic Quantum Theory* (Pergamon Press, New York, 1977).
 - [11] E. G. Adelberger *et al.*, Rev. Mod. Phys. **70**, 1265 (1998).
 - [12] V. M. Galitski, JETP **34**, 104 (1958).